

PEAVINE



Peavine

The squeezed juice from the combined pods, vines, leaves, and stems of the common pea (*Pisum sativum*) is a nutritionally packed source of essential vitamins and a significant source for phenolic compounds. Eating peas and other legumes improves your food quality score (FQS).



Phytoactives

Lignans

Large plant polyphenolic compounds that bypass human digestion, feed gut bacteria, and provide antioxidant activity

Lariciresinol (0.5 mcg/g*)

Pinoresinol (0.07 mcg/g)*

Medioresinol (0.035 mcg/g)*

Syringaresinol (0.04mcg/g)*

Secoisolariciresinol (0.00756 mcg/g)*

Chlorophyll

Green pigment in plants with potential anti-inflammatory, antioxidant, and anti-bacterial activity

Carotenoids

Antioxidants with anti-cancer potential and may lower risk of macular degeneration

Lutein (7.22 mcg/g)**

Zeaxanthin (0.39 mcg/g)**

Flavanols

Promote antioxidant, anticancer, antimicrobial, and anti-inflammatory activity

Catechin (0.1 mcg/g)*

Gallocatechin⁴

Epicatechin (0.1 mcg/g)*

Epigallocatechin⁴

Flavanols

Promote antioxidant activity and promote vascular health

Quercetin²

Kaempferol²

Phenolic Acid

Phytoactive compounds that promote antioxidant activity and promote vascular health

Sinapoyl-glucoside²


Saponins

Support the immune system and promote healthy cholesterol and blood glucose levels

Soyasaponin I³

Soyasaponin Bg³

What is the Whole Food Matrix?



Supports balance immune modulation for healthy inflammation response.

Supports the gut microflora and a healthy metabolic fingerprint of the gut.

Benefits of nutrients food matrix enhances bioavailability by up to 60%.

Organic and adaptive regenerative farming techniques delivers nutrient dense source of key phytonutrients and helps balance healthy lifestyles.

Increased intake of vegetables and fruits in whole food nutrition influences individual epigenetic expression of our health potential.




Gallic Acid Equivalence

What is GAE?

GAE, or “gallic acid equivalence,” indicates levels of important phytoactives available in the plant and extracts. GAE is derived by comparing to the gallic acid reference standard, a simple phenolic substance. Studies have shown that phytoactives in plants contribute to their beneficial effect on development of chronic diseases.

Total Phenolic Concentration

Measured: Total Phenolics as Gallic Acid Equivalence (mg/g)



* Data is mean values from Phenol-Explorer Database¹

** Data on file with WholisticMatters

Values subject to change based on strain and experimental methods

Key Nutrients

Percentages shown as %DV per 5g of dry peavine plant extract

Vitamin K

Vital for blood clotting and healthy bones.

16%

Vitamin E

A micronutrient with antioxidant activity that supports the immune system and metabolism.

13%

Biotin (Vitamin B7)

B vitamin necessary for energy metabolism, histone modification, gene regulation, and cell signaling.

11%

Riboflavin (Vitamin B2)

Water-soluble vitamin vital for energy production, cell function, metabolism, and growth/development.

10%

Magnesium

An essential mineral that supports nerve and muscle function, the immune system, and a healthy heart.

7%

Other Nutrients

(in order of %DV per 5g peavine juice extract)

Copper

Vitamin B6 (Pyridoxal

5'-phosphate)

Iron

Folate

Calcium

Selenium

Niacin (Vitamin B3)

Manganese

Potassium

Protein

Pantothenic acid (Vitamin B5)

Zinc

Phosphorus

Choline

Carbohydrates

Fiber

Thiamin (Vitamin B1)

Lipids



We are dedicated to advancing the latest insights and information available in nutrition therapy and clinical nutrition and to presenting only the most balanced, credible, and reliable clinical nutrition and science available.

WholisticMatters.com

©2019 Standard Process Inc.

References

- Rothwell, J.A., et al., Phenol-Explorer 3.0: a major update of the Phenol-Explorer database to incorporate data on the effects of food processing on polyphenol content. Database, 2013. 2013: p. bat070-bat070.
- Neugart, S., Rohn, S., & Schreiner, M. (2015). Identification of complex, naturally occurring flavonoid glycosides in Vicia faba and Pisum sativum leaves by HPLC-DAD-ESI-MSn and the genotypic effect on their flavonoid profile. Food Research International, 76, 114-121. doi:https://doi.org/10.1016/j.foodres.2015.02.021
- Reim, V., & Rohn, S. (2015). Characterization of saponins in peas (Pisum sativum L.) by HPTLC coupled to mass spectrometry and a hemolysis assay. Food Research International, 76, 3-10. doi:https://doi.org/10.1016/j.foodres.2014.06.043
- Jin, A., Ozga, J. A., Lopes-Lutz, D., Schieber, A., & Reinecke, D. M. (2012). Characterization of proanthocyanidins in pea (Pisum sativum L.), lentil (Lens culinaris L.), and faba bean (Vicia faba L.) seeds. Food Research International, 46(2), 528-535. doi:https://doi.org/10.1016/j.foodres.2011.11.018